

Contribution of Agricultural Practices on Enzymatic Activity in Anopheles Gambiae Populations from Cotton Areas in Northern Benin

Anges YADOLETON*, Jean-Robert KLOTOE, Christophe CHABI

Abstract— To have an idea on the contribution of agricultural practices on biochemical characterization of insecticide resistance in *Anopheles gambiae* populations, *Anopheles* larvae were collected from cotton field from July to September 2017 and reared until F1 generation.

Enzymatic mechanisms (Glutathione-S-Transferase, esterase, monooxygenase P450 and total protein) were investigated from the F1 (the parental population after breeding) populations of *An. gambiae* s.l.

Results from this study showed a significant high level of GST and monooxygenase P450 activities from the wild populations of *An. gambiae* from Banikoara compared to susceptible Kisumu strain ($P>0.05$). However, there is no significant difference in the level activity of Esterase (α and β -Naphthyl) from the wild populations of *An. gambiae* from compared to the susceptible Kisumu strain ($P>0.05$).

These findings confirmed the contribution of agricultural practices on the selection of enzymatic activity in *An. gambiae* populations particularly on Glutathione-S-Transferase, and monooxygenase P450. However, the same level of the wild populations in esterase showed compared to the susceptible Kisumu strain showed that there is no clear link between the use of insecticide and enzymatic activity in *An. gambiae* populations.

Index Terms— *Anopheles gambiae*; Enzyme; Cotton; Vegetable farming; Benin..

I. INTRODUCTION

The scaling up of Long Lasting Insecticidal Nets (LLINs), the extent Indoor Residual Spraying (IRS) and treatment with artemisinin-based combination therapy (ACT) are currently the major international strategies to control malaria, particularly in sub-Saharan Africa [1]. Currently only one insecticide class, the pyrethroids, is commonly used to treat LLINs; pyrethroids have the required dual properties of low mammalian toxicity and rapid insecticidal activity [2], and their repellent or contact irritant effects may enhance the personal protection of LLINs. Unfortunately, resistance to pyrethroids is now widespread in African malaria vectors [3].

In fact, the development of pyrethroid resistance in the primary malaria vectors, *An. gambiae* s.l and *An. funestus* [4] is a serious concern. In the last decade, the emergence of resistance in populations of *An. gambiae* to common classes

Anges YADOLETON, 1Centre de Recherches Entomologiques de Cotonou, Cotonou, Benin

Ecole Normale Supérieure de Natitingou ; Université Nationale des Sciences, Technologies, Ingénierie et Mathématiques (UNSTIM) d'Abomey

Jean-Robert KLOTOE, Ecole Normale Supérieure de Natitingou ; Université Nationale des Sciences, Technologies, Ingénierie et Mathématiques (UNSTIM) d'Abomey

Christophe CHABI, Faculté d'Agronomie. Université de Parakou-Benin

* Corresponding author

of insecticides used in public health has been reported in many African countries with a high allelic frequency of knock down resistance (Kdr) [5-8].

In Benin, with pyrethroid resistance spreading in *Anopheles gambiae*, the national malaria Control Programme in Benin has introduced a carbamate (bendiocarb) as an alternative to pyrethroids for the Indoor Residual Spraying (IRS) [9]. However, resistance to carbamate particularly to bendiocarb has been documented by Aikpon et al [8] three years later after it introducing in areas where IRS strategies were implemented against mosquitoes bites. In fact, a study conducted in Benin, revealed the use of bendiocarb in cotton fields against cotton pests [10].

Benin is still an important producer of cotton in West Africa where 90% of pesticide products are directed against cotton pests [11]. These insecticides are essentially composed of pyrethroids (PYs), organophosphates (OPs) and carbamates which are also the main classes used in public health. The majority of cotton farms observed in northern Benin are located in the upland landscape while the lowland covers the major mosquito breeding sites. According to Akogbeto et al. [12] in Benin, insecticide treatments against cotton pests are applied twice a month, for a timeframe of three consecutive months (between July and October) each year. These treatment periods coincide with the rainy season and correspond to the period of high mosquito densities.

While insecticide resistance associated with *knock down resistance (kdr)* and *Acetylcholinesterase (aceIR)* is well studied at the physiological, behavioural and population level [10; 13- 15] much less is known about the enzymes associated with metabolic resistance.

The present study aimed to provided information on biochemical analysis on *An. gambiae* populations from Cotton field to detect potential increase in mixed function oxidases (MFO), non-specific esterases (NSE) and glutathione S-transferases (GST) activity.

II. MATERIAL AND METHOD

A. Study area

Cotton field at Banikoara located in the north-east of Benin (Figure 1) was chosen for investigation. Cotton production is the main activity of the populations in this district. Cotton crop protection represents 90% of the insecticide use in this district for crops protection [10]. The control strategies implemented against cotton pest especially *Helicoverpa armigera* required a regular repeated applications of insecticides during the cotton plant growing cycle. As recommended by the Institut National des Recherches Agronomiques du Benin (INRAB), six consecutive



Contribution of Agricultural Practices on Enzymatic Activity in Anopheles Gambiae Populations from Cotton Areas in Northern Benin

treatments are applied at two weeks interval to protect the crop against bollworms, leafworms and sucking pests. These insecticides are essentially composed of pyrethroids (PYs), organophosphates (OPs).

B. Mosquito collection

Larvae of *An. gambiae* s.s. were collected at the two sites (Figure 1) and reared at CREC (Centre de Recherche Entomologique de Cotonou) insectary for emergence. Emerging adult females mosquitoes (F_0) aged 6-8 days were fed with guinea pig blood and reared for emergence (F_1) where adults 3-5 days were kept at -80 degrees for biochemical analysis.

C. Biochemical analysis

60 adult females of the wild populations of *An. gambiae* s.s. from the study site (Figure 1) were kept at -80 degrees and were subjected to biochemical based on the methods described by Penilla al [16] to compare the levels of activity of mixed function oxidases (MFO), non-specific esterases (NSE) using α -naphthyl acetate as a substrate and glutathione S-transferases (GST) to the laboratory Kisumu susceptible reference strain.

Individual mosquitoes were homogenized in 200 μ l ml distilled water. Each of 10 ml of the homogenate was used for monooxygenase, glutathion S-transferase and protein assay. The other twenty μ l ml of homogenate was used for esterases assay.




Figure 1: Map of Benin showing the study site

D. Glutathione -S-transferase (GST) assay

10 μ l of each homogenate was transferred to a microplate well followed by 200 μ l of the GSH/CDNB working solution which was prepared by adding 0.060g of glutathione solution(GSH) in 20 ml of Phospahte sodium buffer 0.1M and 0.013gr (in 1 ml of methanol) 1-chloro-2,4-dinitrobenzene (CDNB). The plates were read after 5 mins with the ELISA plate reader at a wave length of 340 nM. GST activity was expressed as: mMoles/ min / mg protein

E. Monooxygenase (Cytochrome p450) assay

10 μ l of homogenate were placed in separate of microtitre plate followed by addition of 80 μ l 0.625M potassium phosphate buffer (pH 7.2). Ten mg of 3,3,5'5', Tetramethyl Benzidine(TMBZ) in 5 ml methanol were prepared and a 15 ml of 0.25 M sodium acetate buffer(pH 5.0) was prepared. Two hundred μ l of the above TMBZ solution was added in to each well followed by 25 μ l of 3% hydrogen peroxide. The plate was read after 2 hours at 630 nm

F. Esterase assay

20 μ l of homogenated were placed in separate wells of microtitre plate. 200 μ l of 0.3 mM Alpha/Beta napthyl

acetate were added to each well. Leave the plate at room temperature for 1 min and then added 50 μ l of fast garnet. After 30 minutes, enzyme activity was determined as an OD value by microplate reader at 450 nm.

G. Protein assay

The total protein content of individual mosquitoes was determined using the Bio -Rad Protein Assay Kit (Bio -Rad Laboratories) in order to detect the differences in size among individuals that might require correction factors for the enzyme assays

H. Data analysis

Biochemical assay data (enzymatic activity per mg protein, levels of MFO, NSE and GST between Kisumu and field populations *An. gambiae* s.s.) were compared using Mann-Whitney non-parametric U-test (Statistica software).

III. RESULTS

Results from biochemical assay showed a significantly higher level of GST activity of GST activity from the wild populations of *An. gambiae* from the study site compared to susceptible Kisumu strain (Mann-Whitney test, P>0.05) (Figure 2). The same trend was observed with Monooxygenase (Cytochrome p450). (Figure 3)




Figure 2 : Glutathione activity of *Anopheles gambiae* populations from the study areas

Contribution of Agricultural Practices on Enzymatic Activity in *Anopheles gambiae* Populations from Cotton Areas in Northern Benin





Figure 3: Mixed Function Oxidases activity of *Anopheles gambiae* populations from the study sites

However, there is no significant difference in the level activity of Esterase (α and β -Naphthyl) from the wild populations of *An. gambiae* from the study site compared to the susceptible Kisumu strain ($P>0,05$)






Figure 4: Esterase activity of *Anopheles gambiae* populations from the study site

insecticides for pest control.

V. CONCLUSION

These findings confirmed the contribution of agricultural practices on the selection of enzymatic activity in *An. gambiae* populations particularly on Glutathione-S-Transferase, and monooxygenase P450. However, the same level of the wild populations in esterase showed compared to the susceptible Kisumu strain showed that there is no clear link between the use of insecticide and enzymatic activity in *An. gambiae* populations.

COMPETING INTERESTS

The authors declare that they have no competing interests.

REFERENCES

- [1] World Health Organization Global Malaria Programme. Conditions for use of long-lasting insecticidal nets treated with a pyrethroid and piperonyl butoxide. December 2015.
- [2] Zaim M, Aitio A, Nakashima N. Safety of pyrethroid-treated mosquito nets. *Medical and Veterinary Entomology* 2000;14(1):1-5.
- [3] Ranson H, Lissenden N. Insecticide resistance in African *Anopheles* mosquitoes: a worsening situation that needs urgent action to maintain malaria control. *Trends in Parasitology* 2016;32(3):187-96
- [4] N'Guessan R, Corbel V, Akogbeto M, Rowland M. Reduced efficacy of insecticide-treated nets and indoor residual spraying for malaria control in pyrethroid resistance area, Benin. *Emerging Infectious Diseases* 2007;13(2):199-206

Contribution of Agricultural Practices on Enzymatic Activity in Anopheles Gambiae Populations from Cotton Areas in Northern Benin

- [5] Corbel V, N'Guessan R, Brengues C, Chandre F, Djogbenou L, Martin T, Akogbeto M, Hougard JM, Rowland M: Multiple insecticide resistance mechanisms in *Anopheles gambiae* and *Culex quinquefasciatus* from Benin, West Africa. *Acta Trop* 2007, 101:207-216.
- [6] Yadouleton AW, Asidi A, Djouaka RF, Braïma J, Agossou CD, Akogbeto MC: Development of vegetable farming: a cause of the emergence of insecticide resistance in populations of *Anopheles gambiae* in urban areas of Benin. *Malar J* 2009, 8:103.
- [7] Czeher C, Labbo R, Arzika I, Duchemin JB: Evidence of increasing Leu-Phe knockdown resistance mutation in *Anopheles gambiae* from Niger following a nationwide long-lasting insecticide-treated nets implementation. *Malar J* 2008, 7:189.
- [8] Aïkpon R, Agossa F, Ossè R, Oussou O, Aizoun N, Oké-Agbo F, et al. Bendiocarb resistance in *Anopheles gambiae* s.l. populations from Atacora department in Benin, West Africa: a threat for malaria vector control. *Parasit Vectors*. 2013;6:192
- [9] Akogbéto M, Padonou GG, Gbénou D, Irish S, Yadouleton A. Bendiocarb, a potential alternative against pyrethroid resistant *Anopheles gambiae* in Benin, West Africa. *Malar J*. 2010, 9: 204
- [10] Yadouleton AW, Martin T, Padonou G, Chandre F, Alex A, Djogbenou L, Dabiré R, Aïkpon R, Glitho I, Akogbeto MC: Cotton pest management strategies on the selection of pyrethroid resistance in *Anopheles gambiae* populations in northern Benin. *Parasites and Vectors* 2011, 4:60
- [11] IFDC. L'état du marché des intrants agricoles au Bénin. *Technical Bulletin*. 2005, 83 pp
- [12] Akogbeto M C, Djouaka RF, and Kinde-Gazard DA: Screening of pesticide residues in soil and water samples from agricultural settings. *Malar J* 2006, 5: 22
- [13] Santolamazza F, Mancini E, Simard F, Qi Y, Tu Z, della Torre A Insertion polymorphisms of SINE200 retrotransposons within speciation islands of *Anopheles gambiae* molecular forms. *Malar J*. 2008;7:163
- [14] Lalji S, Ngondi JM, Thawer NG, Tembo A, Mandike R, Mohamed A, et al. School distribution as keep-up strategy to maintain universal coverage of long-lasting insecticidal nets: implementation and results of a program in Southern Tanzania. *Glob Health Sci Pract*. 2016;4:251–263
- [15] David JP, Ismail HM, Chandor-Proust A, Paine MJ. Role of cytochrome P450 s in insecticide resistance: impact on the control of mosquito-borne diseases and use of insecticides on Earth. *Philos Trans R Soc Lond B Biol Sci*. 2013;368:20120429
- [16] Penilla, R.P., Rodriguez, A.D., Hemingway, J., Torres, J.L., Arredondo-Jimenez, J.I. & Rodriguez, M.H. (1998) Resistance management strategies in malaria vector mosquito control. Baseline data for a large scale field trial against *Anopheles albimanus* in Mexico. *Medical and Veterinary Entomology* 12, 217–233
- [17] Joncour G: Lutte anti-palustre au Dahomey Rapport no 13: Ministère de la Santé Publique 1959.